



The effects of Chloroquine on the histology and stereology of Sprague - Dawley rats heart

Akinribido F.A, Noronha C.C, Okanlawon O.A (2011)

Dept Department of Biological Sciences, Bells University of Technology Ota, Ogun State

Dept Of Anatomy, College of Medicine, University of Lagos, Idi-Araba, Lagos State

(Research Work)

Correspondence to: Akinribido F.A, fumibos@yahoo.co.uk; +2348023654975

ABSTRACT

Chloroquine has been reported to have a depressant effects on cardiac muscle and chloroquine like quinine, reduces excitability and conductivity of cardiac muscle (Taylor and Francis 1987).it is therefore necessary to study the effects of chloroquine on the heart .

This study was designed to determine the effects of administration of chloroquine on the heart. Ten male rats were exposed to chloroquine injection intra peritoneally/IP) for three days. The treated rats received 0.125ml/100g body weight of chloroquine phosphate injection intraperitoneally. Control rats received the same amount of normal saline intraperitoneally.

The histology of the chloroquine treated heart was compared with the controls. It was observed that chloroquine caused malformations this tissue. Investigation confirmed defects in microscopic structures. There were few blood vessels in the treated rats compared with controls and there were constrictions in the structures of treated rats compared with controls.

Stereologically, the estimated absolute volume $V=Vv(\text{structure}) \times V(\text{ref})$ was determined and compared chloroquine caused a reduction in the absolute volume of the heart blood vessels (coronary artery) compared with controls

Keywords: chloroquine, heart, Sprague Dawley rats, histology, stereology.

INTRODUCTION

Approximately 50-70% of chloroquine in plasma is bound to plasma proteins. The tissues exhibit particularly high binding to chloroquine especially those containing melanin, for example the retina. Significant binding also occurs in the liver, kidney and spleen. Chloroquine (Resochin, Avloclor, Nivaquine, Arelen) C₁₈H₂₆ClN₃ 7- Chloro -4- (4'- diethylamino-1'-methylamino) quinoline. Chloroquine is a white powder with a bitter taste, prepared by chemical synthesis .it is available as sulphate and phosphate salts. The sulphate (1 in 3) and the phosphate (1 in 4) are soluble in water. Chloroquine is best known as an antimalarial agent but it is also used in the treatment of rheumatoid arthritis. Chloroquine is effective against the erythrocytic stages of all four plasmodium species which cause human malaria with the exception of matured plasmodium falciparum gametocytes. The exact mechanisms of the action of chloroquine against malaria parasites are not fully understood. Parasitized red cells accumulate approximately 100-600 times as much chloroquine. The concentration of chloroquine in malaria parasite requires energy and is thought to require a membrane. There are three theories on the way state as that chloroquine, being a basic compound, is protonated in the lysosomes thus raising lysosomal pH. This effect may raise the intralysosomal pH above a critical level all bring about loss lysosomal function. This would reduce the parasite's digestion of heamoglobin, and thus prevent its growth. Chloroquine intercalates into double stranded DNA and inhibits both DNA and RNA synthesis. The intercalation theory suggests that chloroquine may be bound with increased affinity by certain parts of the genome and be toxic to the malaria parasite by selective accumulation in specific genes, inhibiting their expression. The ferriprotophyrin IX (FP) which inhibits sequestration of FP into malaria pigment. This could impair heamoglobin degradation and permits damage to the food vacuole sufficient to discharge its Ph gradient. antimalaria activity is possessed equally by the enantiomers of chloroquine and the main metabolite desethylchloroquine is also active against chloroquine- sensitive Plasmodium. Chloroquine also has anti-inflammatory activity. The concentrations of chloroquine or hydrochloroquine found in serum in the treatment

of rheumatoid disease raise the pH of acid vesicles in mammalian cell within 3-5 min in vitro. This and the observation that the view that chloroquine and hydroxychloroquine act in the rheumatic disease by raising the pH of acid vesicles. Effects of raised vesicle pH include inhibition lysosomal proteolysis, interference with the targeting of acid proteases and inhibition of cellular maturation .raise pH in the macrophage vesicle can interfere with antigen processing. This is thought to be the explanation for the impaired antibody response to pre-exposure to human diploid cell rabies vaccine found in individual receiving concurrent chemoprophylaxis with chloroquine. In addition, chloroquine inhibits the chemotactic response of mononuclear cells and suppresses lymphocytes transformation.

MATERIALS AND METHODS

The twenty male Sprague- Dawley rats were collected from the Animal House of the College of Medicine University of Lagos Akoka, Lagos State.

They weighed between 100-150g and were fed with the normal rat feed from Pfizer PLC Ikeja Lagos. Weight of animals was taken twice daily throughout the duration of the experiment. Ten female rats were used as controls. The remaining ten female rats were labelled by ear puncture as treated rats and kept in cages. Administration of drug was 0.125ml of chloroquine /100g body weight for 3 days intraperitoneally. Chloroquine phosphate injection was obtained from the community pharmacy of the Lagos university teaching hospital (40mg/ml chloroquine phosphate injection).The control received the same quantity of normal saline.

Animal Sacrifice

At the expiration of the treatment the animals were sacrificed by diethyl ether decapitation and the heart was removed for morphological and histological assessment.

Histological Analysis

The twenty male rats were sacrificed as discussed earlier after treatment with the chloroquine phosphate injection .The heart was removed and fixed in Bouin's fluid. The specimen of equal length was cut transversely and longitudinally into serial cross sections of 3µm normal thickness with Reichert Jung Supercut Microtome for control and treated rats. The tissue was sectioned using tissues preparation tissues method with heamatoxylin and eosin stains and examined the light binocular microscope at a magnification of 100 and 400 respectively.

Stereological Analysis

The vertical sections of the histochemical preparation of stratum length of 0.5cm from 10 control and 10 treated rats heart was made at a final print magnification of 100 and 400 respectively.

5 slides will be obtained from the control and 5 slides from the treated rats.

For each of the fractions, the $N_T/A \equiv$ number of tests points counted on rats hearts coronary artery per unit area of the fractions were estimated by point counting method using the forbidden rule Hans Gundersen,1977) which states that any structure that touches the forbidden line must not be counted. The reference volume $V(\text{ref})$ of heart coronary artery was estimated by point counting (Wiebel, 1979, Gundersen et al, 1988).

At Magnification (M) = 100 final magnification using a Square Grid of test point diameter (d) =1.2cm apart. The test system used in the light microscopic analysis within a square frame measuring 20cm x 20cm onto which microscopic image was projected using a wild leitz microscope equipped with a mirror at a magnification of 25 on a white screen.

Estimated $V(\text{ref}) = (\text{stratum length}) \times \frac{d^2}{M^2} \times \text{mean } N/A \text{ (structure)}.$

d= diameter of test grid

M=magnification of projection

The relevant volume density of heart coronary artery of the fractions V_v (structure) were estimated on the same section at a final magnification of 100. Each field was projected onto a test system consisting of three sets of points with numerical densities in the ratio 1:4:16. The corresponding distance between the test points of each set were 4.8, 2.4 and 1.2cm respectively.

The criteria for test point design and allocation were based on efficiency considerations; thus approximately the same number of test points (which does not need to exceed 200) should be in each structure within each organ (Gundersen and Jensen, 1987; Gundersen et al., 1988; Cruz Orive and Wiebel; 1990). The required volume density of the fractions were estimated as follows:

$$\text{Estimated } V_v(\text{structure}) = N_{vR} \times N_T/A(\text{structure})$$

V_v = volume density N_{vR} = numerical density ratio

Finally, the absolute volume of heart coronary artery was estimated using this equation.

$$V(\text{structure}) = V_v(\text{structure}) \times V(\text{ref})$$

$V(\text{structure})$ = Absolute volumes of structure

$V_v(\text{ref})$ = Reference volume of structure

STATISTICS

Statistical analysis was carried out using t- Distribution (t- test).

RESULTS

TABLE1: MEAN NUMBER OF TEST POINTS PER UNIT AREA (N_T/A) CORONARY ARTERY

GROUP n=20	Mean (N_T/A)
CONTROL (CO)	140
CHLOROQUINE TREATED (CQ)	135

CO=CONTROL RATS

CQ= CHLOROQUINE TREATED RATS

TABLE2: MEAN ESTIMATED ABSOLUTE VOLUMES (CM^3) OF CORONARY ARTERY

TISSUE	CONTROL RATS n=10	CHLOROQUINE TREATED RATS n=10
Heart(coronary artery)	$3.53 \times 10^{-1} \pm 0.86^a$	$3.28 \times 10^{-1} \pm 1.63^b$

a=Mean±S.E.M

b=p<0.05

DISCUSSION

Chloroquine caused defects in the structures of the heart. They were few blood vessels in the treated rats compared with controls. There were also constrictions in this structure. Stereologically, chloroquine caused a reduction in the heart coronary artery absolute volume compared with controls.

CONCLUSION

The heart is very important as it circulates blood round the body precautions should be taken as to when chloroquine should be taken under the supervision of a licensed medical practitioner

REFERENCES

1. Bray PG, Ward SA, O'Neill PM. Plasmodium falciparum malaria: rosettes are disrupted by quinine, artemisinin, mefloquine, primaquine, pyrimethamine, chloroquine and proguanil. *Curr Top Microbiol Immunol.* 2005;295:3-38.
2. Chawira AN, Warhurst DC. Noculum effect leads to overestimation of in vitro resistance for artemisinin derivatives and standard antimalarials: a Gambian field study. *J Trop Med Hyg.* 1987 Feb;90(1):1-8.
3. Cooper RA, Carucci DJ. *Curr Opin Investig Drugs.* 2007 Feb;8(2):117-24.
4. Duraisingh MT, Jones P, Sambou I, von Seidlein L, Pinder M, Warhurst DC. Drug-resistant malaria in Bangladesh: an in vitro assessment. *Parasitology.* 1999 Nov;119 (Pt 5):435-40.
5. Ehrhardt S, Eggelte TA, Kaiser S, Adjei L, Burchard GD, Anemana SD, Bienzle U, Mockenhaupt FP. In vitro stage-specific sensitivity of Plasmodium falciparum to quinine and artemisinin drugs. *Med Trop (Mars).* 1998;58(3 Suppl):13-7
6. Hastings IM, Bray PG, Ward SA. Analysis of pfcrt, pfmdr1, dhfr, and dhps mutations and drug sensitivities in Plasmodium falciparum isolates from patients in Vietnam before and after treatment with artemisinin. *Science.* 2002 Oct 4;298(5591):74-5.
7. Khan MA, Smego RA Jr, Razi ST, Beg MA. Emerging drug-resistance and guidelines for treatment of malaria. *J Coll Physicians Surg Pak.* 2004 May;14(5):319-24.
8. Lelièvre J, Berry A, Benoit-Vical F. Artemisinin and chloroquine: do mode of action and mechanism of resistance involve the same protagonists?
9. Majori G. Developing artemisinin based drug combinations for the treatment of drug resistant falciparum malaria: A review. *Parassitologia.* 2004 Jun;46(1-2):85-7.
10. Meshnick SR. Large-scale surveillance of Plasmodium falciparum crt(K76T) in northern Ghana.
11. Ngo T, Duraisingh M, Reed M, Hipgrave D, Biggs B, Cowman AF. Rational use of drugs against Plasmodium falciparum. *Am J Trop Med Hyg.* 2003 Mar;68(3):350-6.
12. Noedl H, Faiz MA, Yunus EB, Rahman MR, Hossain MA, Samad R, Miller RS, Pang LW, Wongsrichanalai C. Artemisinin antimalarials: mechanisms of action and resistance. *Am J Trop Med Hyg.* 2003 Feb;68(2):140-2.
13. Noedl H, Wernsdorfer WH, Krudsood S, Wilairatana P, Viriyavejakul P, Kollaritsch H, Wiedermann G, Looareesuwan S. Antimalarial drug resistance and combination chemotherapy. *Am J Trop Med Hyg.* 2001 Dec;65(6):696-9.
14. Olliaro PL, Taylor WR. [Drug resistant tropical malaria in the Republic of Guinea (West Africa)]. *J Postgrad Med.* 2004 Jan-Mar;50(1):40-4
15. *Philos Trans R Soc Lond B Biol Sci.* 1999 Apr 29;354(1384):739-49.
16. Popov AF, Popova NI, Chirkov VP, Kamara F. Proteomic approaches to studying drug targets and resistance in Plasmodium. *Med Parazitol (Mosk).* 2003 Apr-Jun;(2):41-3.
17. Skinner TS, Manning LS, Johnston WA, Davis TM. *Antimicrob Agents Chemother.* 2007 Sep;51(9):3407-9. Epub 2007 Jun 11.
18. The effect of artemisinin combined with standard antimalarials against chloroquine-sensitive and chloroquine-resistant strains of Plasmodium falciparum in vitro. *Curr Drug Targets Infect Disord.* 2004 Mar;4(1):41-51.
19. Trotta RF, Brown ML, Terrell JC, Geyer JA. Defective DNA repair as a potential mechanism for the rapid development of drug resistance in Plasmodium falciparum. *Int J Parasitol.* 1996 May;26(5):519-25
20. Warhurst DC, Duraisingh MT. In vivo-in vitro model for the assessment of clinically relevant antimalarial cross-resistance. *Trans R Soc Trop Med Hyg.* 2001 Jul-Aug;95(4):345-6.
21. White N. Quinolines and artemisinin: chemistry, biology and history. *Biochemistry.* 2004 May 4;43(17):4885-91.